

PACS: 11.10.-z

ON INFLUENCE OF INTENSIVE STATIONARY ELECTROMAGNETIC FIELD ON THE BEHAVIOR OF FERMIONIC SYSTEMS

V.N. Rodionov, A.M. Mandel, E.V. Arbuzova

General Physics Department,

Moscow State Geological Prospecting Academy,

117997, Miklukho -Maklaya, 23

Abstract.

Exact solutions of Schrödinger and Pauli equations for charged particles in an external stationary electromagnetic field of an arbitrary configuration are constructed. Green functions of scalar and spinor particles are calculated in this field. The corresponding equations for complex energy of particles bounded by short range potential are deduced. Boundary condition typical for δ - potential is not used in the treatment. Explicit analytical expressions are given for the shift and width of a quasistationary level for different configurations of the external field. The critical value of electric field in which the idea of quasistationary level becomes meaningless is calculated. It is shown that the common view on the stabilizing role of magnetic field concerns only scalar particles.

I. Introduction. Formulation of the problem.

The influence of external electromagnetic fields on nonrelativistic reactions with charged particles and on the behavior of bound systems (like atoms, ions and atomic nucleuses) has being investigated systematically for a long time see, (for example, [1–15]). However, a set of problems, in our opinion, requires additional studying. First, it is the appearance of physically meaningless divergences in equations for the complex energy of quasistationary bound states. Second, the question about the behavior of fermionic systems in super strong electromagnetic fields is not enough elucidated. Note that the basic results in this field of physics were received by nonperturbative methods. Usually the exact solutions of Schrödinger equations with Hamiltonians taking into account the presence of external fields are used [16, 17]. It is important, that the field can become intensive not only due to increasing of the strength, but also due to decreasing of binding energy of the system. In other words, in the vicinity of the reaction threshold an arbitrary external field is strong. Third, the common view on the

stabilizing role of magnetic field in decays of quasistationary states is inadequate for spinor particles [18,19]. That view is caused by the fact that spinor states of electrons usually are not taken into accounts in nonrelativistic reactions. In this paper we treat a part of these deficiencies.

Let us consider a charged particle bounded by a short range potential like a δ -potential and located in an external stationary electromagnetic field of an arbitrary configuration. Note that the potential of zero-radius is a rather widely spread in the literature approximation for a multi-electronic atom field and especially for a negative ion field, and also for the field of nuclear forces [12,20]. In the general case the external field is defined by three independent parameters: strengths of the magnetic H and electric E fields and the angle ϕ between them.

The process of particle coming out of a δ -well leads to the transition from an isolated energy level to a level in a continuous spectrum, *i.e.* to a decay of the bound state. The external electromagnetic field influences on this transition in two ways. First, it forms wave functions of the “free” state and, second, shifts and broads the bound level in a δ -well. As the result the initial state of the particle in the external field becomes quasistationary. The most adequate instrument for the investigation of such states is the complex energy formalism (for example, [4,12,21])

$$W = W_0 + \Delta W - i\Gamma/2, \quad (1)$$

where $W_0 < 0$ is the energy of the nonperturbative bound level, ΔW is the real shift caused by the external field, and Γ is the width of a level associated with the decay probability of the bound state. It is important that a separate consideration of the level shift and its width is possible only in a weak external field. In an intensive field it is necessary to investigate these quantities in common.

The main purpose of this article is to derive an equation for the complex energy in a stationary external field by the *method of an analytical continuation* and to solve it for some field configurations. This treatment is principally different from the traditional derivation using the boundary condition typical for the δ -well [4,14,15].

The general structure of the paper and its main results are as follows. In the second Section explicit solutions for Schrödinger and Pauli equations are constructed both for scalar and spinor particles in an external stationary electromagnetic field. Then Green functions of the particles are constructed with its help. In spite of rather complicated configuration of the external field, these functions can be represented in terms of elementary functions. That substantially simplifies their further applications.

In the third Section the expression for the bound level width in the external field is deduced with help of the Green functions. Then the equation for the complex energy

of the bound particle in the δ -well is built by the method of analytical continuation. It has several features different from the ones of the equation traditionally used. First, our equation is not transcendental and it is explicitly resolved with respect to the complex energy. That is why its right handed side (one fold integral) does not contain exponential divergences typical for equations used earlier. Thus it does not require any additional regularization. Second, we consistently take into account the spin states, which is fundamentally important in the presence of magnetic field.

Finally in the fourth Section the equation for the complex energy is solved in several cases of physical interest. It is well known that the electric field destroys the bound state of a charged particle in a potential well. Thus we calculated the critical value of the strength of the electric field in which the width of the level becomes comparable with its depth in the potential well. In such a field the notion of an isolated level loses meaning [21]. When the strengths of the electric and magnetic fields are comparable the latter one gives only small corrections to the usual effects: the Stark shift and the broadening due to the tunnel effect. But the signs of these corrections are different for scalar and spinor particles. In the case when the magnetic field is dominating the behaviors of the complex energy of the mentioned particles are principally different. For scalar particles the magnetic field exponentially decreases the bound level width and therefore stabilizes it. It was assumed that the same effect takes place also for electrons (e.g. [5, 10, 14, 15]). But as was shown by the authors of this paper in [18, 19], the strong magnetic field linearly increases the level width for particles with spin 1/2. Hence in the presence of an external magnetic field it is necessary to take into account the spin of the given particle and its contribution to the complex energy of the quasistationary level even in non-relativistic problems.

II. The wave function and the Green function of a charged particle in an external stationary electromagnetic field

In order to receive the wave function of a free charged spinless particle in a stationary electromagnetic field of a general configuration, it is necessary to solve the Schrödinger equation with the following Hamiltonian:

$$\hat{H}_{Sh} = -\frac{\hbar^2}{2m}\Delta - i\hbar\omega_H x \frac{\partial}{\partial y} + \frac{1}{2}m\omega_H^2 x^2 + eEx \sin \varphi + eEz \cos \varphi, \quad (2)$$

where e and m are the charge and the mass of the particle; E and H are the strengths of the electric and the magnetic fields, respectively; φ is the angle between the corresponding vectors; $\omega_H = eH/mc$ is the cyclotron frequency. We suppose that the

magnetic field has the orientation along z axis, and the vector of the electric field strength lies in the xz plane.

The Schrödinger equation with the Hamiltonian (2) can be solved by the standard method of variables separation. The moving of the particle in a plane normal to the magnetic field has the same character as the one in the absence of electric field. Corresponding eigenfunctions are Hermit functions, each of them is related to a certain Landau level. In this case the coordinate x_0 of the orbit centre depends on the conserved transverse momentum p_y and on the transverse component of the electric field $E \cos \varphi$. The movement of the particle along z axis is similar to the one in the uniform electric field, and it can be described by Airy functions [21]. It is not difficult to see that the wave function of the charged particle in the field of the considered configuration has the form

$$\psi_n(\vec{r}, t) = N u_n(\rho) B(\xi) \exp \left[\frac{i}{\hbar} (y p_y - t \tilde{W}) \right], \quad (3)$$

where N is a normalization coefficient, the Hermit function $u_n(\rho)$ satisfies the equation

$$u_n''(\rho) + (2n + 1 - \rho)u_n(\rho) = 0$$

where n is the number of the Landau level, and the argument of this function is

$$\rho(x) = \left(\frac{m\omega_H}{\hbar} \right)^{1/2} \left[x + \frac{p_y}{m\omega_H} - \frac{eE \sin \varphi}{m\omega_H^2} \right].$$

Function $B(\xi)$ in (3) is proportional to the Airy function Ai regular at infinity

$$B(\xi) = \frac{(2m)^{1/3}}{\pi (eE \cos \varphi)^{1/6} \hbar^{2/3}} Ai(\xi)$$

with the argument

$$\xi(z) = \left(\frac{2meE \cos \varphi}{\hbar^2} \right)^{1/3} (z - z_0),$$

where z_0 is defined by the total energy W of the particle in the external field

$$\tilde{W} = \hbar\omega_H \left(n + \frac{1}{2} \right) + eE z_0 \cos \varphi. \quad (4)$$

The solution of the Schrödinger equation constructed in that way is valid for a scalar particle. In order to take into account spin states of the particle in the concerned non-relativistic approximation it is necessary to pass on to the Pauli equation (see, for example, [21]) for the spinor wave function $\bar{\psi}_{n\sigma}(\vec{r}, t)$

$$i\hbar \frac{\partial}{\partial t} \bar{\psi}_{n\sigma}(\vec{r}, t) = \hat{H}_P \bar{\psi}_{n\sigma}(\vec{r}, t),$$

where the Hamiltonian of the Pauli equation involves the energy of the spin interaction with the external stationary magnetic field $\vec{H}$:

$$\hat{H}_P = \hat{H}_{Sh} + \frac{e\hbar}{2mc} \vec{\sigma} \vec{H},$$

and $\vec{\sigma}$ are the Pauli matrixes. Taking into account the concrete orientation of the magnetic field this Hamiltonian is reduced to the matrix form

$$\hat{H}_P = \begin{pmatrix} \hat{H}_{Sh} + \hbar\omega_H/2 & 0 \\ 0 & \hat{H}_{Sh} - \hbar\omega_H/2 \end{pmatrix}.$$

If we put now into the Pauli equation the spinor wave function of the mixed state with an arbitrary time dependence of the coefficients

$$\bar{\psi}_{n\sigma}(\vec{r}, t) = \psi_n(\vec{r}, t) \begin{pmatrix} a_1(t) \exp(-i\omega_H t/2) \\ a_2(t) \exp(i\omega_H t/2) \end{pmatrix}, \quad (5)$$

where the scalar function $\psi_n(\vec{r}, t)$ is defined by the formula (3), we get

$$da_1/dt = da_2/dt = 0.$$

It means, first, that the spin projection on the z axis is conserved. This conservation is provided by the homogeneity and the stationarity of the magnetic field. Second, the energy of the particle is defined, instead of the expression (4), by the more common formula

$$\tilde{W} = \hbar\omega_H(n + 1/2 + \sigma) + eEz_0 \cos \varphi, \quad (6)$$

where $\sigma = 0$ for the scalar particle and $\sigma = \pm 1/2$ for the spinor one.

Thus four constants (p_y, z_0, n, σ) unambiguously parameterize the wave function of the particle with spin 1/2 in the external field under consideration.

Let's construct now a time depending Green function of a charged particle in the given field. With the purpose of some simplification of calculations we, without limiting the generality, suppose one of arguments of point-to-point Green function to be zero. The function will be a diagonal matrix of a kind

$$\widehat{G}(\vec{r}, t; \vec{0}, 0) = \sum_{n=0}^{\infty} \int_{-\infty}^{\infty} dz_0 \int_{-\infty}^{\infty} dp_y G_{Sh}(n, z_0, p_y) \begin{pmatrix} \exp(-i\omega_H t/2) & 0 \\ 0 & \exp(i\omega_H t/2) \end{pmatrix} \quad (7)$$

where the scalar Green function $G_{Sh}(n, z_0, p_y)$ corresponds to a Schrödinger equation

$$G_{Sh}(n, z_0, p_y) \sim u_n(x)u_n(0)Ai(z)Ai(0) \exp\left[\frac{i}{\hbar}(p_y y - t\tilde{W})\right].$$

The integral over z_0 included in (7) is easily calculated using the integral representation for the Airy function. The integral over p_y with Hermit functions is similar to the one that was considered in Ref. [22]. The series in Lagerre polynomials formed after that is summarized over n with help of the algorithm described in [23] (see also [24]). As a result the scalar part of the Green function in a stationary electromagnetic field is expressed in terms of elementary functions:

$$G_{Sh}(\vec{r}, t; \vec{0}, 0) = \left(\frac{m}{2\pi i}\right)^{3/2} \frac{\omega_H}{2t^{1/2}} \sin^{-1}\left(\frac{\omega_H t}{2}\right) \exp\left(\frac{iS}{\hbar}\right), \quad (8)$$

where

$$S = \frac{mz^2}{2t} - \frac{eEzt}{2} \cos \varphi - \frac{(eE \cos \varphi)^2 t^3}{24m} + \frac{m\omega_H}{4} \left[(x^2 + y^2) \cot\left(\frac{\omega_H t}{2}\right) - 2xy \right] \\ + \frac{1}{2} eE x t \sin \varphi + \frac{eE \sin \varphi}{\omega_H} \left[\frac{\omega_H t}{2} \cot\left(\frac{\omega_H t}{2}\right) - 1 \right] \left(y + \frac{eEt \sin \varphi}{2m\omega_H} \right). \quad (9)$$

If we assign in (8), (9) the mass of the particle to unity and perform obvious changes in notation we exactly reproduce formulas (A.7) - (A.9) from Ref. [14]. At $\varphi = \pi/2$ the relations (8), (9) describe the Green function of a scalar particle in the crossed field [9, 10, 24].

3. Derivation of the equation for the complex energy of a bound particle in a stationary external field by the method of analytical continuation

The expression for the probability P of the decay of a particle bound state per unit time (or for the width Γ of the bound level) in the considered field can be received with help of the Green function, using the procedure developed in Refs. [11, 25]. This probability has the form

$$\begin{aligned}
P &= \pi \left(\frac{2}{m} \right)^{3/2} \left(\frac{|W_0|}{\hbar} \right)^{1/2} \int_{-\infty}^{\infty} dt G_{Sh}(\vec{0}, t; \vec{0}, 0) f_{\sigma} \left(\frac{\omega_H t}{2} \right) \exp \left(\frac{i}{\hbar} W_0 t \right) \\
&= \frac{\omega_H}{2} \left(\frac{|W_0|}{\pi \hbar} \right)^{1/2} \exp \left(-i \frac{3\pi}{4} \right) \int_{-\infty}^{\infty} \frac{dt}{t^{1/2}} \frac{f_{\sigma}(\omega_H t/2)}{\sin(\omega_H t/2)} \exp \left[\frac{i}{\hbar} (S + W_0 t) \right], \quad (10)
\end{aligned}$$

where W_0 is the energy of the unperturbed bound level (1), and S at zero coordinates is defined by the formula

$$S = -\frac{(eE \cos \varphi)^2 t^3}{24m} + \frac{(eE \sin \varphi)^2 t}{2m\omega_H^2} \left[\frac{\omega_H t}{2} \cot \left(\frac{\omega_H t}{2} \right) - 1 \right]. \quad (11)$$

Poles of the integrand function in points $t_n = 2\pi n/\omega_H$ are passed from below.

Function $f_{\sigma}(a)$ is identically equal to unity for scalar particles, but for spinors it depends on the polarization of the bound particles in the initial state and of free particles in the final one. In the general case

$$f_{\sigma}(\omega_H t/2) = a_1^2 \exp(-i\omega_H t/2) + a_2^2 \exp(i\omega_H t/2), \quad (12)$$

where $a_1(a_2)$ are the amplitudes of the probability introduced in (5), that the spin of a particle is directed along (or against) the magnetic field. The normalization condition demands that $a_1^2 + a_2^2 = 1$. For example, the most realistic situation in the process of ionization is the one where bound electrons are not polarized and the detector does not distinguish between polarizations of free electrons. In this case for the calculation of the total probability of ionization it is necessary to average over polarizations of the initial particles and to sum over polarizations of the final ones (see, for example, [21]). Then $a_1^2 = a_2^2 = 1/2$ and $f_{\sigma} = \cos(\omega_H t/2)$.

As it is known the width of the bound level is proportional to the probability of its decay $\Gamma = \hbar P$. For analytical continuation let us rewrite (10) in such a way that the integration in it is carried out on the positive semi-axis. With this purpose it is necessary to resolve the exponent in (10) into the real and the imaginary parts and in the integral contained $\cos[(S + W_0 t)/\hbar]$ to bypass the zero point from below on an infinitesimal contour. Such a bypass cancels the divergence proportional to $t^{-1/2}$ of the above mentioned integral at zero. As a result the formula for the level width takes the form

$$\frac{\Gamma}{4|W_0|^{1/2}} = \frac{1}{2} \left(\frac{\hbar}{\pi} \right)^{1/2} \int_0^{\infty} \frac{dt}{t^{3/2}}.$$

$$\left\{ \frac{\omega_H t}{2 \sin(\omega_H t/2)} \sum_{k=1,2} a_k^2 \sin \left[\frac{1}{\hbar} \left(S + W_0 t + (-1)^k \frac{\hbar \omega_H}{2} t \right) - \frac{\pi}{4} \right] + \sin \frac{\pi}{4} \right\}. \quad (13)$$

It becomes obvious now that the right hand side of (14) represents the imaginary part of the complex expression

$$(-W)^{1/2} = \frac{1}{2} \left(\frac{\hbar}{\pi} \right)^{1/2} \int_0^\infty \frac{dt}{t^{3/2}} \cdot \left\{ \frac{\omega_H t}{2 \sin(\omega_H t/2)} \sum_{k=1,2} a_k^2 \exp \left[\frac{i}{\hbar} \left(S + W_0 t + (-1)^k \frac{\hbar \omega_H}{2} t \right) - \frac{i\pi}{4} \right] - \exp \left(-\frac{i\pi}{4} \right) \right\} \quad (14)$$

and the left hand side is nothing else than the imaginary part of the expansion (1)

$$(-W)^{1/2} = (-W_0 - \Delta W + i\Gamma/2)^{1/2} \approx |W_0|^{1/2} - \frac{\Delta W}{2|W_0|^{1/2}} + i \frac{\Gamma}{4|W_0|^{1/2}}.$$

It is logical to assume that the real part of (14) determines the level shift due to the external field. If we use now the identical transformation (recalling that $W_0 < 0$)

$$\int_0^\infty \frac{dt}{t^{3/2}} \left[\exp \left(i \frac{t W_0}{\hbar} \right) - 1 \right] = -2 \left(\frac{\pi |W_0|}{\hbar} \right)^{1/2} \exp \left(\frac{i\pi}{4} \right), \quad (15)$$

we receive the final equation for the complex energy of a bound level in the stationary external field of a general configuration

$$(-W)^{1/2} - (-W_0)^{1/2} = \frac{1}{2} \left(\frac{\hbar}{i\pi} \right)^{1/2} \int_0^\infty \frac{dt}{t^{3/2}} \exp \left(\frac{i}{\hbar} W_0 t \right) \cdot \left\{ \frac{\omega_H t}{2} \frac{f_\sigma(\omega_H t/2)}{\sin(\omega_H t/2)} \exp \left(\frac{iS}{\hbar} \right) - 1 \right\}, \quad (16)$$

where S is defined by formula (11).

We see that the received equation is resolved explicitly with respect to W . Note, that all integrals in intermediate expressions used in the derivation as well as in the final formula (16) are *finite*. Besides the condition of the relative smallness of the initial level width and shift has been used. It imposes certain restrictions on the area of applicability of (16). However, such restrictions are incorporated in the concept of a

quasistationary energy level [21]. More strict mathematical derivation of the equation similar to (16) is given in [27].

In appendix A of paper [14] a more commonly used variant of derivation of the equation for the complex energy in the considered configuration of an external field (see also [4, 5, 10, 15, 20]) is described. Taking into account the spins of particles, we can get the equation

$$(-W)^{1/2} - (-W_0)^{1/2} = \frac{1}{2} \left(\frac{\hbar}{i\pi} \right)^{1/2} \int_0^\infty \frac{dt}{t^{3/2}} \exp \left(\frac{i}{\hbar} W t \right) \cdot \left\{ \frac{\omega_H t}{2} \frac{f_\sigma(\omega_H t/2)}{\sin(\omega_H t/2)} \exp \left(\frac{iS}{\hbar} \right) - 1 \right\}, \quad (17)$$

which differs from the equation (16) only by the substitution $W_0 \rightarrow W$ in its right hand side. Thus the two approaches leading to slightly different equations are formulated for the description of quasistationary systems bounded by short-range forces. In this connection it is useful to compare the areas of their applicability and to understand how much are different their solutions.

First note that usually an iteration method is used to solve equation (17). Therefore the substitution $W_0 \rightarrow W$ is accepted on the first step of the iterative procedure, reducing thereby (17) to (16). Therefore the received equation (16) is treated usually as *an approximate one*, working only in the case of weak external fields. Thus, the choice of one of the alternative approaches suggested above is practically reduced to the question: whether iterative corrections of higher orders are necessary or it is needed to stop on the “first iteration”.

Obviously, the equation (17) for the complex energy is transcendental and, hence, more complicated than (16). Solving a more complicated equation is meaningful only if it gives a more exact or an essentially new result. Therefore it is important to discuss approximations used by the derivation of the specified relations. At the derivation of equation (16) we used conditions $\Gamma \ll |W_0|, |\Delta W| \ll |W_0|$, which impose certain restrictions from above on values of external fields. However, the presence of such restrictions follows both from the non-relativistic character of the problem, and from the initially assumed presence of “the shallow bounded level” in a deep narrow potential well [12]. Moreover, the concept of a quasistationary level assumes the smallness of its decay probability [21]. We shall discuss the last circumstance in detail below.

However, it is not possible at all to estimate restrictions on the values of external fields, admitted at the derivation of (17). The matter is that all intermediate integrals (formulas (A.6), (A.13) - (A.19) in Ref. [14]) as well as the final integral in Eq. (17)

are exponentially divergent . At that it is emphasized in works [4,5,20] that the typical for a δ -well boundary condition is valid only for a complex value of W

$$G(\vec{r}, \vec{0}; W) \underset{r \rightarrow 0}{\approx} \frac{m\hbar^{1/2}}{2\pi r} - \frac{m^{3/2}}{(2\hbar)^{1/2}\pi} (-W)^{-1/2} + \tag{18}$$

$$i^{-1/2} \left(\frac{m}{2\pi}\right)^{3/2} \int_0^\infty \frac{dt}{t^{3/2}} \exp\left(i\frac{Wt}{\hbar}\right) \left\{ \frac{\omega_H t}{2} \sin^{-1}\left(\frac{\omega_H t}{2}\right) \exp\left(\frac{iS}{\hbar}\right) - 1 \right\} + O(r).$$

Therefore it remains not clear to terms of which order it is necessary to relate the last integral at $ImW < 0$.

Finally, in our opinion, the key condition

$$G(\vec{r}, \vec{0}; W) \underset{r \rightarrow 0}{\approx} G_0(\vec{r}, \vec{0}; W_0) \underset{r \rightarrow 0}{\approx} \frac{m\hbar^{1/2}}{2\pi r} - \frac{m^{3/2}}{(2\hbar)^{1/2}\pi} (-W_0)^{-1/2} + O(r). \tag{19}$$

on which total procedure of a derivation [14] is based causes some doubt. Namely the comparison of asymptotics (18) and (19) gives the traditional equation for a complex energy (17). The condition (19) actually demands that not only the singular part but also the finite one of the asymptotic expansion of the Green function at zero does not depend on external fields. As far as we know, arguments of physical character for the benefit of this were not given in the literature. Nevertheless the asymptotic of the field Green function at zero (18) contains in the *finite* part the integral, divergent for complex values of W . In these conditions the requirement of the equality of finite terms is an obvious excess of the accuracy of the used approximation. This will be shown in the following section for the case of a pure electric external field.

The mentioned above difficulties of the derivation of expression (17) are characteristic not only for an approach suggested in Ref. [14]. They are typical for any derivation of equation (18) using the boundary condition of the δ -potential with a complex energy. There are no such problems in Refs. [4,5,10] for the reason that their authors determine the complex energy, in contrast to (1), with a positive imaginary part. It removes the problem of divergences in formulas like (17)-(19), but contradicts to the classical definition of the complex energy. In agreement with such a definition all probabilities in a quasistationary state should exponentially increase in time [21].

Thus, it does not seem to be possible, in our opinion, to derive equation (17) with the complex energy W in the right hand side consistently and accurately from the mathematical point of view. It is especially problematic to trace all approximations used in such a derivation.

4. Solving equations for the complex energy of a bound particle for some configurations of external fields.

Let us stop now on some variants of the solution of the received equations for simple configurations of an external field. It is stated in Ref. [28] that the elimination of divergences in integrals like (18) by the substitution $W \rightarrow W_0$ is incorrect. Let's discuss further a rather simple configuration of an external electrostatic field from the point of view of the validity and the applicability of equations (16) and (17). We use the notation of Ref. [28]: the energy ε and the strength of the electrostatic field F are measured in units of $|W_0|$ and in characteristic atom units, respectively:

$$\varepsilon = \frac{W}{|W_0|}, \quad F = \frac{eE\hbar}{(2m|W_0|^3)^{1/2}}.$$

Of course, the particle spin does not show itself in an electrostatic external field. At that our equation (16) takes the form [27]

$$(-\varepsilon)^{1/2} - 1 = (-\varepsilon_0)^{1/2} + \frac{F^{1/3}}{\pi} \left[Ai' (Bi' + iAi') + \varepsilon_0 F^{-2/3} Ai (Bi + iAi) \right], \quad (20)$$

where the common argument of the Airy functions Ai, Bi and of their derivatives Ai', Bi' is $-\varepsilon_0 F^{-2/3}$, and the zero approximation gives $\varepsilon_0 = -1$. Formula (17), as it is affirmed in [15, 28], gives in such an external field the transcendental equation

$$1 + \frac{1}{\pi} F^{1/3} \left[Ai' (Bi' + iAi') + \varepsilon F^{-2/3} Ai (Bi + iAi) \right] = 0. \quad (21)$$

and the complex energy is contained in arguments of the Airy functions $-\varepsilon F^{-2/3}$. However, note once again that it is impossible to pass from (17) to (21) by means of identical transformations. The Airy functions on the left handed side of equation (21) are finite while the integral on the right hand side of (17) diverges. It is easy to see, at what stage of transformations the infinity “can disappear”. For example, it is possible to pass from the general expression (17) to (21) with help of transformation (15) with real W_0 . However, the integral on the left hand side of (15) diverges explicitly for complex W with a negative imaginary part, but the right handed side remains finite. The mentioned in the previous section excess of the accuracy in the derivation of equation (17) from the comparison of asymptotics (18) and (19) reveals here.

At the same time, it is possible to receive the transcendental Eq. (21) by analytical continuation of the right handed side of (20) over the argument of the Airy functions

into the complex plane, replacing ε_0 by complex ε . As the calculations show, values of ε determined from (21) are not differ too much from the values that gives the explicit formula (20). In a weak field ($F \ll 1$) both relations reproduce the well-known asymptotic expression [4]

$$-\varepsilon = 1 + \frac{F^2}{16} + \frac{i}{4}F \exp\left(-\frac{4}{3F}\right),$$

if to be limited to first non-vanishing terms of the Airy functions expansions. In the strong field $F = 1$ (i.e. comparable with intra-atomic one) it is easy to get from (20)

$$-\varepsilon \approx 1,0411 + 0,0451 \cdot i$$

and the numerical solution of (21) gives [15, 28]

$$-\varepsilon \approx 1,0442 + 0,0388 \cdot i.$$

Besides, there is a critical value of the electric field F_{cr} at which $Re(-\varepsilon) = Im(-\varepsilon)$, i.e. the energy gap between the “shifted and broadened” level and the continuous spectrum disappears. Obviously, that usual concept of a quasistationary level from [21] loses the meaning for such intensive electric fields and requires an additional definition. In accordance with (20), that critical value is $F_{cr} \approx 13,26$. The transcendental equation (21) gives $F_{cr} \approx 11,38$.

Finelly note that there is a well-known analogy between the complex energy in a theory of quasistationary system decays and the complex dielectric permeability in physics of semiconductors (for example, [7, 8, 29, 30]). Equations for the complex dielectric permeability are derived, as a rule, on the basis of well developed dispersive methods.

One more configuration of an external field which we consider is essentially important to show the role of the magnetic field. For simplicity we put the angle $\varphi = 0$ and average over polarizations of bound particles. It is convenient to measure the magnetic field like the electric one in natural atom units (see, for example, [14]):

$$h = \frac{eH\hbar}{mc|W_0|}.$$

If $h \leq F \ll 1$ it is possible to receive [19] from our equation (16)

$$(-\varepsilon)^{1/2} - 1 = \frac{F^2}{32} + \frac{h^2}{48a} + i\frac{F}{8} \left(1 + \frac{h}{3aF^2}\right) \exp\left(-\frac{4b^{3/2}}{3F}\right).$$

Here the dependent on spin factors a and b take values $a = -2$, $b = 1 + h/2$ for a scalar particle and values $a = b = 1$ for a spinor one. Therefore, the corrections to the complex energy due to a magnetic field essentially depend on spins of particles. The magnetic field decreases the width of a scalar particle bound level and by that stabilizes it. On the contrary, the width of the level of a particle with spin $1/2$ increases in a magnetic field, i.e. the latter strengthens the destabilizing action of an electric field. That difference is still more noticeable in a strong magnetic field [18]. This field suppresses exponentially the withdrawal of scalar particles from the δ -well, but intensifies it linearly for particles with spin:

$$Im(-\varepsilon)_{\sigma=0}^{1/2} \sim h^{1/2} \exp\left(-\frac{2^{1/2}h^{3/2}}{3F}\right); \quad Im(-\varepsilon)_{\sigma=1/2}^{1/2} \sim \frac{h}{F^{1/3}} Ai^2(F^{-3/2}).$$

If we distinguish particle polarizations after ionization it is not difficult to see with help of equation (16) and condition (12) that electrons with spins directed along a magnetic field ($a_1^2 = 1/2$, $a_2^2 = 0$) in the initially non-polarized beam behave like scalar particles. Thus, the increasing of the level width in a magnetic field is provided only by electrons with spins oriented against the field.

The reason for the phenomenon described above can be seen from equation (6) for the total energy of an electron in an external field. Energies of Landau levels enhance with the increase of the magnetic field. That complicates tunneling of a particle from the bound state. Only one exception is the ground state of an electron with spin directed against the magnetic field. The energy of this state does not depend on H at all, but its contribution to the total probability of ionization increases with the increase of the magnetic field.

Therefore for an adequate description of the influence of a magnetic field on the behavior of fermionic quasistationary systems it is necessary to take into account spin states of fermions even if the considered problem is not relativistic.

References

- [1] Keldysh L. V. 1958 *Zh. Exsp. Teor. Fiz.* **34**. 1138 (Engl. transl. 1959 *Sov. Phys. - JEPT.* **34**. 788)
- [2] Franz W. 1958 *Z. Naturf. A.* **13**. 484
- [3] Zel'dovich Ya. B. 1961 *Zh. Exsp. Teor. Fiz.* **39**. 776 (Engl. transl. 1961 *Sov. Phys. - JEPT* **12**. 542)

- [4] Demkov Yu. N. and Drukarev G. F. 1964 *Zh. Exsp. Teor. Fiz.* **47**. 918 (Engl. transl. 1965 *Sov. Phys. - JEPT* **20**. 614)
- [5] Demkov Yu. N. and Drukarev G. F. 1965 *Zh. Exsp. Teor. Fiz.* **49**. 257 (Engl. transl. 1965 *Sov. Phys. - JEPT* **22**. 182)
- [6] Keldysh L. V. 1964 *Zh. Exsp. Teor. Fiz.* **47**. 1945 (Engl. transl. 1965 *Sov. Phys. - JEPT* **20**. 1037)
- [7] Seraphin B. O. and Bottka N. 1965 *Phys. Rev.* **139**. A560
- [8] Aspnes D. E. 1966 *Phys. Rev.* **147**. 554
- [9] Ritus V. I. 1966 *Zh. Exsp. Teor. Fiz.* **51**. 1544 (Engl. transl. 1967 *Sov. Phys. - JEPT* **25**. 1027)
- [10] Drukarev G. F. and Monozon B. S. 1971 *Zh. Exsp. Teor. Fiz.* **61**. 956 (Engl. transl. 1972 *Sov. Phys. - JEPT* **34**. 509)
- [11] Nikishov A. I. 1972 *Zh. Exsp. Teor. Fiz.* **62**. 562 (Engl. transl. 1972 *Sov. Phys. - JEPT* **35**. 298)
- [12] Baz A. I, Zel'dovich Ya. B. and Perelomov A. M. 1971 *Scattering, Reactions and Decays in Nonrelativistic Quantum Mechanics* 2nd edn (Moscow: Nauka) (in Russian) (Engl. transl. - 1st edn, US Department of Commerce, Natural Technical Information Service, N69. 25848)
- [13] Zel'dovich Ya. B, Manakov N. L. and Rapoport L. P. 1975 *Usp. Fiz. Nauk* **117**. 569 (Engl. transl. 1975 *Sov. Phys.- Usp.* **18**. 920)
- [14] Popov V. S, Karnakov B. M. and Mur V. D. 1998 *Zh. Exsp. Teor. Fiz.* **113**. 1579 (Engl. transl. 1998 *Sov. Phys. - JEPT* **86**. 860)
- [15] Manakov N. L, Frolov M. V, Starace A. F. and Fabrikant I. I. 2000 *J Phys B: At Mol Opt Phys* **33**. R141
- [16] Dodonov V. V, Man'ko V. I. and Nikonov D. E. 1992 *Phys. Lett. A* **162**. 359
- [17] Kleber M. 1994 *Phys. Rep.* **236**. 331
- [18] Rodionov V. N, Kravtsova G. A. and Mandel A. M. 2002 *Zh. Exsp. Teor. Fiz. Lett.* **75**. 435 (Engl. transl. 2002 *Sov. Phys. - JEPT Letters* **75**. 363)

- [19] Rodionov V. N, Kravtsova G. A. and Mandel A. M. 2003 *Zh. Exsp. Teor. Fiz. Lett.* **78**. 253 (Engl. transl. 2003 *Sov. Phys. - JEPT Letters* **78**. 218)
- [20] Demkov Yu. N. and Ostrovsky V. N. 1988 *Zero-Range Potentials and Their Applications in Atomic Physics* (New York: Plenum)
- [21] Landau L. D. and Lifshitz E. M. 1992 *Quantum Mechanics* 4th edn (Oxford: Pergamon)
- [22] Klepikov N. P. 1954 *Zh. Exsp. Teor. Fiz.* **26**. 19
- [23] Rodionov V. N. 1998 *Zh. Exsp. Teor. Fiz.* **113**. 21 (Engl. transl. 1998 *Sov. Phys. - JEPT* **86**. 11)
- [24] Feynman R. P. and Hibbs A. R. 1965 *Quantum Mechanics and Path Integral* (New York: McGraw-Hill Book Company)
- [25] Rodionov V. N. 1997 *Zh. Exsp. Teor. Fiz.* **111**. 3 (Engl. transl. 1998 *Sov. Phys. - JEPT* **84**. 1)
- [26] Nikishov A. I. and Ritus V. I. 1983 *Zh. Exsp. Teor. Fiz.* **85**. 1544 (Engl. transl. 1983 *Sov. Phys. - JEPT* **58**. 14)
- [27] Kadyshevskii V.G. and Rodionov V. N. 2000 *Theor. Math. Phys.* **125**. 1668 (Engl. transl. 2000 *Theor. Math. Phys.* **125**. 1654)
- [28] Manakov N. L, Frolov M. V, Borca B. and Starace A. F. 2000 *Zh. Exsp. Teor. Fiz. Lett.* **72**. 426 (Engl. transl. 2000 *Sov. Phys. - JEPT Letters* **52**. 294)
- [29] Roth L. M, Lax B. and Zwerdling S. 1959 *Phys. Rev.* **114**. 90
- [30] Nussenzveig H. M. 1972 *Causality and dispersion relations* (New York; London: Acad. Press)